

Planck-scale effects on WIMP dark matter

M. S. Boucenna, R. A. Lineros*, and J.W.F. Valle

*AHEP Group, Instituto de Física Corpuscular – C.S.I.C./Universitat de València
Edificio de Institutos de Paterna, Apartado 22085, E-46071 València, Spain*

(Dated: March 2, 2013)

Weakly Interactive Massive Particles are among the most motivated candidates invoked to solve the Dark Matter puzzle. The stability of Dark Matter (DM) typically results from a global protective symmetry. However, the presence of gravitational effects may cause its explicit breaking. We study the presence of such effects on WIMPs and show that in addition to providing a new source of signal for indirect DM searches and an interplay with direct detection, a large class of decaying Dark Matter models can be embedded into the WIMP paradigm. Taking into account the recent limits on DM lifetimes, we show that forbidding only the dimension five (Planck-suppressed) operators is generally sufficient to ensure the validity of the WIMP candidate.

PACS numbers: 95.35.+d, 98.80.Cq

It has been conjectured and argued that non-perturbative gravitational effects break all global (continuous and discrete alike) symmetries [1, 2]. The original motivation was based on black hole physics arguments but since then perturbative string theory has confirmed this conjecture in various cases [2]. Such gravitational effects have already been considered in a number of scenarios and models invoked to solve fundamental theoretical problems [3–11]. Here, we turn our attention to the consequence of such a claim on dark matter phenomenology, in the particular case of weakly interacting dark matter candidates.

There is, indeed, considerable evidence in favor of the existence of dark non-baryonic matter in the universe; arising from very different scales. However, despite the ongoing efforts since decades, identifying the nature of DM remains a major problem in modern physics.

Weakly Interacting Massive Particles (WIMPs) [12] present in various extensions of the Standard Model (SM) are among the best motivated candidates for cold DM, as they quite naturally account for the observed relic abundance. Indeed, their abundance at freeze-out epoch is:

$$\Omega_{\text{CDM}} h^2 \simeq 0.1 \frac{3 \times 10^{-26} \text{ cm}^3 \text{s}^{-1}}{\langle \sigma v \rangle_{\text{f.o.}}}, \quad (1)$$

where $\langle \sigma v \rangle_{\text{f.o.}}$ is the thermal average of the annihilation cross-section of DM into lighter particles at freeze-out. For masses in the range from GeV to few TeV, WIMPs have a typical weak strength interactions cross-sections, passing all constraints arising from Big-Bang Nucleosynthesis (BBN), Cosmic Microwave Background (CMB) and structure formation. This non-trivial interplay between the weak-scale physics and the cosmological dark matter has been dubbed as the “WIMP miracle”.

Besides theoretical motivation and consistency, the WIMP paradigm provides positive direct and indirect detection prospects, owing to the eventual recoil off nuclei and annihilation into photons, neutrinos and/or charged particles. The increasing level of sensitivity that WIMP direct [13–15] and indirect detection experiments [16–18] are reaching and the possibility of complementary searches [19] with the LHC open great hopes that this goal will finally be reached soon.

However, WIMPs are generally *assumed* to be stable particles. This is achieved in most models by means of an imposed global symmetry (usually a Z_2) that forbids the decay of the DM candidate to lighter states. Therefore unless the symmetry responsible for the DM stability is local, it is expected to be broken at short distances by gravitational effects leading to a decaying WIMP DM. We can, however, arrange the model such that there is an accidental symmetry stabilizing the DM at the level of renormalizable operators, leaving the planck-mass suppressed dimension five and higher operators responsible of the decay.

From the phenomenological point of view absolute stability is not a necessary condition for DM candidates. Instead, what is required is a lifetime larger than the age of the Universe $H^{-1} \approx 10^{17} \text{s}$, where H is the Hubble constant. Cosmic and gamma rays analysis constrain the lifetime of a DM candidate even further to be $\tau_{\text{DM}} > 10^{26} \text{s}$ [20–23]. Therefore the main question to be addressed is what are the constraints on the gravity-induced decay to preserve the validity of the DM candidate.

On the other hand, decaying DM candidates are commonly assumed to have a non-thermal origin, hence not to be WIMPs. Which is indeed the case of particles with masses below or much above the weak scale (keV-majorons [8, 24, 25], sterile neutrinos [26], Superheavy DM [27]) and/or with extremely weak couplings with SM particles (gravitinos [28, 29] for e.g.), however, this not necessarily the case of weak-scale DM particles.

*Multidark Fellow.

In this Letter we show that a number of phenomenologically interesting decaying DM candidates with electroweak scale masses and couplings fit within a WIMP scenario, providing a well controlled cosmological history (thermal production) and direct detection possibilities. Alternatively the indirect detection prospects of WIMPs are enhanced by the new signals which arise from the decay, in particular in regions where annihilations alone deliver a very faint one.

Simplest model: In order to illustrate the discussion, we consider a very simple model, exhibiting generic WIMP dark matter features in a broad phenomenological range. The SM is extended by S_i ($i = 1 \dots N$) real gauge-singlets under an imposed parity symmetry to which the SM itself is blind. Note that the stabilizing symmetry need not be a Z_2 ; any global (discrete or continuous) symmetry that forbids the decay of the lightest S_i is a valid choice. Alternative SM extensions providing viable DM candidates with positive detection prospects have been considered for instance in refs. [30–32].

The above scenario provides a production and thermalization mechanism via Higgs boson exchange (the so-called Higgs portal [33]). For small enough the mass splitting [34], co-annihilations exist between the LSP (lightest singlet particle), which is the DM candidate S_1 , and the remaining $S_{i>1}$. In a conventional WIMP dark matter scenario our LSP would be stable and the scalar potential would read as (summation over indexes is understood):

$$\mathcal{V}_{\text{sym}} = -\mu_h^2 H^\dagger H + \mu_{ij}^2 S_i S_j + \lambda_{ij} S_i S_j H^\dagger H + \lambda'_{ijkl} S_i S_j S_k S_l. \quad (2)$$

For the sake of simplicity, we consider the case $N = 2$ and $\mu_{ii}^2 > 0$. The latter precludes spontaneous breaking of the Z_2 symmetry, and the associated domain walls.

The dark matter phenomenology of the model is then essentially determined by 4 parameters [48]: M_{S_1} , M_{S_2} , λ_{11} and λ_{12} . In the absence of coannihilations ($\lambda_{12} \rightarrow 0$), λ_{11} is responsible of the DM abundance as well as direct (σ_{SI}) and indirect detection ($\langle \sigma_A v \rangle$) cross-sections. The strength of coannihilations is controlled by λ_{12} , while λ_{22} is the Higgs coupling of S_2 (the next-to-LSP) to the Higgs scalar which has direct impact on the relic abundance of S_1 . This happens in regions of the parameter space where the mass splitting is small and coannihilations and annihilations are inefficient to reproduce a good relic abundance. In this case the latter could arise partly from the early decays of $S_2 \rightarrow S_1$.

We present in Fig. 1 the variation of the thermal average of the annihilation cross-section times velocity at present time, $\langle \sigma_A v \rangle$, compatible with DM relic abundance fixing the Higgs mass to 126 GeV [35, 36] and varying the couplings within the limits of perturbativity. The results were obtained using the `Micromegas` code [37, 38].




Figure 1: Annihilation cross section $\langle \sigma_A v \rangle$ compatible with DM relic abundance versus DM mass. No inhibited annihilation is shown in black, inhibited annihilations (see text) at freeze-out larger than 90 and 99 % are shown by the green and red shaded regions, respectively. The thermal cross section $\langle \sigma v \rangle = 3 \times 10^{-26} \text{cm}^2/\text{s}$ and FERMI’s constraints for annihilation into $b\bar{b}$ is also shown.

In the presence of pure annihilations, $\langle \sigma_A v \rangle$ reproduces the thermal value $\langle \sigma v \rangle_{\text{f.o.}} \simeq 3 \times 10^{-26} \text{cm}^3/\text{s}$ (Eq. 1).

Deviations from the thermal value exist in regions where the cross-section is velocity-dependent as in the case of the Breit–Wigner enhancement at the threshold of the Higgs pole [39]. The expected indirect detection signal in this case is at the edge of FERMI’s [40] sensitivity. Another big deviation occurs at the Higgs boson mass and is associated with the contact interaction term.

In regions where $\langle \sigma v \rangle_{\text{f.o.}}$ is dominated by coannihilation ($S_1 S_2 \rightarrow \text{SM} + \text{SM}$ and/or $S_2 S_2 \rightarrow \text{SM} + \text{SM}$) processes, the annihilation cross-section gets inhibited and drops well below the expected thermal value. These “inhibited annihilation” regions can be achieved in various ways. Coannihilation is just a possible mechanism, common to many DM models (e.g. MSSM, 2HDM), that we use solely for illustration purposes.

Inhibition mechanisms are often a necessary ingredient in order to get or extend an allowed region in the parameters space of the considered model [41, 42]. The general result, however, is that the WIMP’s indirect detection signal becomes much fainter.

We will see now that the mere fact of considering the global symmetry responsible of the stability of DM as an approximate one, due to the presence of gravitational effects, leads to new observational signals which allow for potentially detectable WIMP-like decaying dark matter. To this end, we add to $\mathcal{V}_{\text{sym}}$ (Eq. 2) an effective potential that breaks explicitly the stabilizing symmetry through




Figure 2: The η -flux versus DM mass (see text for details). We present estimations for 90%, 99%, and no inhibited annihilation (solid green, red, and black lines resp.). Purple lines correspond to decay signal induced by the operator $\hat{\mathcal{O}}_5^{bbh}$ for fixed κ_5 values. A combined signal annihilation and decay is in red dotted-dashed line assuming $\kappa_5 = 3 \times 10^{-8}$ and 99% inhibited annihilation. Bound from FERMI on annihilation into $b\bar{b}$ [40] is also shown.

non-renormalizable terms suppressed by powers of the Planck mass $M_{pl} \approx 10^{19}$ GeV:

$$\mathcal{V}_{\text{non-sym}} = \sum_{n>4} \frac{\kappa_n}{M_{pl}^{n-4}} \hat{\mathcal{O}}_n, \quad (3)$$

where $\hat{\mathcal{O}}_n$ are operators of dimension n that explicitly break the stabilizing symmetry and κ_n are in general complex parameters. This potential clearly leads to the decay of S_1 . Assuming that these corrections are small enough to keep the considered WIMP a viable DM candidate, i.e. adequately long induced lifetime, then we expect a new genuine signal in regions that were previously dark.

To illustrate the interplay between decay and annihilation we consider the following operators $\hat{\mathcal{O}}_5^{ffh} = Y_f H^\dagger F_L f_R S_1$, where F is an SM fermion, Y_f its Yukawa coupling and κ_5 a free parameter. We parametrize the coupling of these dimension five operators as a scaling factor times a Yukawa as a way to model the gravitational effects without claiming a deeper motivation. These operators lead to a decay lifetime $\tau_{DM} = \hbar/\Gamma^{ff}$ where

$$\Gamma^{ff}(M_{DM}) = \sum_f \frac{N_c}{8\pi^2} \left(\frac{\kappa_5 Y_f v}{M_{pl}} \right)^2 M_{DM} \left(1 - \frac{4m_f^2}{M_{DM}^2} \right)^{\frac{3}{2}}, \quad (4)$$

is the DM decay width into a fermions of mass m_f , v is the vacuum expectation value of the higgs boson and N_c

is the color number (3 for quarks and 1 for leptons). For a 50 GeV DM, we would require κ_5 to be smaller than $\sim 10^{-7}$ in order to fulfill the observational constraints.

In Fig. 2, we compare constraints for gamma-rays flux for DM annihilation and decay assuming that the signal comes only from the production of $b\bar{b}$ through the operator $\hat{\mathcal{O}}_5^{bbh}$.

In order to compare the decay and annihilation signals we define the η -flux:

$$\eta(M_{DM}) = \begin{cases} \frac{1}{4\pi} \frac{\langle \sigma_A v \rangle J_{\Delta\Omega}^{\text{ann}}}{2M_{DM}^2} & \text{for annihilations} \\ \frac{1}{4\pi} \frac{J_{\Delta\Omega}^{\text{dec}}}{2\tau_{DM} M_{DM}} & \text{for decays} \end{cases}, \quad (5)$$

where $J_{\Delta\Omega}$ is the angular averaged line of sight integral of the DM density (squared) for decaying (annihilating) WIMP. We consider the $J_{\Delta\Omega}$ reported in Ref. [22] to estimate the fluxes. The η -flux is directly related to the observable gamma-rays flux by:

$$\Phi_\gamma(E, M_{DM}) = \eta(M_{DM}) \times \frac{dn_\gamma(E)}{dE}, \quad (6)$$

where dn_γ/dE is the photon spectrum per single annihilation (or decay) event. This quantity allows us to compare on the same footing both annihilation and decay signals with FERMI constraints. In our particular case, we compare the signal related to the production of $b\bar{b}$ pairs.

We notice that fluxes coming only from inhibited annihilations (same regions as in Fig. 1) lead to signals that are faint and well below observational sensitivities of indirect DM searches. In the mass range from 8 to 100 GeV, inhibited annihilation smaller than 99% exhibit some tension with direct detection constraints (Fig. 3). However, after including the signal originating from the decay modes, for a “reasonable” choice of the κ_n parameters, we can obtain a sizable signal over all the mass range. For heavy states ($> \mathcal{O}(100)$ GeV), FERMI is not sensitive enough to probe the thermal annihilation cross-section and decay modes naturally dominates the signal.

Higher order operators are highly suppressed by the Planck scale and one would require very large couplings ($\sim 10^{10}$) to be able to see a signal from the decay. Thus, the only operators leading to a sizable decay signals are of dimension five. In other words, forbidding dimension five operators is enough to ensure a stable WIMP DM particle in contrast with the dramatic alteration of the axion solution of the strong CP problem by such operators [4, 5].

We now turn to the direct detection signal. This can be calculated in terms of the DM mass and parameterized by σ_{SI} , as shown in Fig. 3. We note that regions with inhibited annihilations are associated with reduced values of σ_{SI} . This model-dependent feature is common




Figure 3: Spin-independent cross section versus DM mass. As in Fig. 1, we present cases where inhibited annihilations produce compatible SI cross section thanks to correlations with annihilation processes. WIMPs with masses larger than 8 GeV would produce extremely faint gamma-ray fluxes in order to fulfill direct detection constraints. Exclusions from XENON100 [43], CDMS [44, 45], and EDELWEISS [46] are shown.

to many constructions where $M_{DM} < M_W$ and annihilations are dominated by quark final states (cf. ref[47] for a particular case). The correlation between σ_{SI} and $\langle\sigma_A v\rangle$ is strong and the inhibition of annihilations is translated as a weakening or absence of direct detection signal. This correlation is certainly important to enlarge the space of parameters allowed within the limits established by XENON100 [43], CDMS [44, 45], and EDELWEISS [46] experiments and opens an interesting interplay between phenomenologically favored light WIMP DM annihilation, decay and direct detection. In other words, models allowing inhibition mechanisms of the WIMP annihilations suffer from the quasi-absence of indirect detection signal. However, taking into account new decay channels reopens the indirect detection potential.

Conclusions: In this Letter, we have considered the effect of explicit gravitational breaking of the global symmetry protecting WIMP DM stability. We have highlighted the importance of taking into account the rich interplay between direct and indirect detection of (decaying) WIMP DM.

In models where inhibited annihilations are present, the expected signal of indirect detection is very faint. However, including the WIMP DM decays induced by the Planck scale effects, provides an extra source of signal making the WIMP DM detectable. On the other hand, a large class of decaying DM candidates, with masses in

the WIMP range and weak-scale couplings, accommodate a WIMP thermal production scenario. Operators of order six and more do not lead to sizable decays, unless very special constructions are considered, and the DM candidate can be seen as stable.

Finally, an unambiguous detection of a mixed state of decay and annihilation signals may offer a very interesting window into Planck scale physics offering an unexpected phenomenological ramification of the WIMP paradigm.

Acknowledgments: We thank N. Fornengo and M. Taoso for their comments and suggestions upon reading the manuscript. This work was supported by the Spanish MEC under grants FPA2011-22975 and MULTIDARK CSD2009-00064 (Consolider-Ingenio 2010 Programme), by Prometeo/2009/091 (Generalitat Valenciana), by the EU ITN UNILHC PITN-GA-2009-237920.

-
- [1] S. B. Giddings and A. Strominger, Nuclear Physics B **307**, 854 (1988).
 - [2] T. Banks and N. Seiberg, PRD **83**, 084019 (2011), 1011.5120.
 - [3] M. Frigerio, T. Hambye, and E. Masso, Physical Review X **1**, 021026 (2011), 1107.4564.
 - [4] M. Kamionkowski and J. March-Russell, Phys.Lett. **B282**, 137 (1992), hep-th/9202003.
 - [5] R. Holman, S. D. Hsu, T. W. Kephart, E. W. Kolb, R. Watkins, et al., Phys.Lett. **B282**, 132 (1992), hep-ph/9203206.
 - [6] E. Masso, F. Rota, and G. Zsembinski, Phys.Rev. **D70**, 115009 (2004), hep-ph/0404289.
 - [7] E. K. Akhmedov, Z. G. Berezhiani, R. N. Mohapatra, and G. Senjanović, Physics Letters B **299**, 90 (1993), hep-ph/9209285.
 - [8] V. Berezhinsky and J. Valle, Phys.Lett. **B318**, 360 (1993), hep-ph/9309214.
 - [9] V. Berezhinsky, A. S. Joshipura, and J. Valle, Phys.Rev. **D57**, 147 (1998), hep-ph/9608307.
 - [10] A. de Gouvêa and J. W. F. Valle, Physics Letters B **501**, 115 (2001), arXiv:hep-ph/0010299.
 - [11] I. Z. Rothstein, K. S. Babu, and D. Seckel, Nuclear Physics B **403**, 725 (1993), arXiv:hep-ph/9301213.
 - [12] G. Jungman, M. Kamionkowski, and K. Griest, Phys.Rept. **267**, 195 (1996), hep-ph/9506380.
 - [13] R. Bernabei, P. Belli, and others., European Physical Journal C **56**, 333 (2008), 0804.2741.
 - [14] C. E. Aalseth, P. S. Barbeau, N. S. Bowden, B. Cabrera-Palmer, J. Colaresi, J. I. Collar, S. Dazeley, P. de Lurgio, J. E. Fast, N. Fields, et al., Physical Review Letters **106**, 131301 (2011), 1002.4703.
 - [15] G. Angloher, M. Bauer, I. Bavykina, A. Bento, C. Bucci, C. Cierniak, G. Deuter, F. von Feilitzsch, D. Hauff, P. Huff, et al., ArXiv e-prints (2011), 1109.0702.
 - [16] D. Hooper and L. Goodenough, Physics Letters B **697**, 412 (2011), 1010.2752.
 - [17] C. Weniger, ArXiv e-prints (2012), 1204.2797.
 - [18] N. Fornengo, R. Lineros, M. Regis, and M. Taoso, Physical Review Letters **107**, 271302 (2011), 1108.0569.
 - [19] G. Bertone, D. G. Cerdeño, M. Fornasa, L. Pieri, R. Ruiz

- de Austri, and R. Trotta, *Phys. Rev. D* **85**, 055014 (2012), 1111.2607.
- [20] A. Ibarra and D. Tran, *JCAP* **2**, 21 (2009), 0811.1555.
 - [21] A. Ibarra, D. Tran, and C. Weniger, *JCAP* **1**, 9 (2010), 0906.1571.
 - [22] X. Huang, G. Vertongen, and C. Weniger, *JCAP* **1**, 42 (2012), 1110.1529.
 - [23] A. Esmaili, A. Ibarra, and O. L. G. Peres, *ArXiv e-prints* (2012), 1205.5281.
 - [24] M. Lattanzi and J. W. F. Valle, *Physical Review Letters* **99**, 121301 (2007), 0705.2406.
 - [25] F. Bazzocchi, M. Lattanzi, S. Riemer-Sorensen, and J. W. F. Valle, *JCAP* **8**, 13 (2008), 0805.2372.
 - [26] S. Dodelson and L. M. Widrow, *Phys.Rev.Lett.* **72**, 17 (1994), hep-ph/9303287.
 - [27] D. J. Chung, E. W. Kolb, and A. Riotto, *Phys.Rev.Lett.* **81**, 4048 (1998), hep-ph/9805473.
 - [28] F. Takayama and M. Yamaguchi, *Physics Letters B* **485**, 388 (2000), arXiv:hep-ph/0005214.
 - [29] D. Restrepo, M. Taoso, J. W. F. Valle, and O. Zapata, *Phys. Rev. D* **85**, 023523 (2012), 1109.0512.
 - [30] C. P. Burgess, M. Pospelov, and T. ter Veldhuis, *Nucl. Phys.* **B619**, 709 (2001), hep-ph/0011335.
 - [31] J. McDonald, *PRD* **50**, 3637 (1994), hep-ph/0702143.
 - [32] Y. Mambrini, *Phys. Rev. D* **84**, 115017 (2011), 1108.0671.
 - [33] B. Patt and F. Wilczek (2006), hep-ph/0605188.
 - [34] K. Griest and D. Seckel, *Phys.Rev.* **D43**, 3191 (1991).
 - [35] ATLAS collaboration, talk by F. Gianotti for the ATLAS collaboration, 13/12/2011.
 - [36] CMS collaboration, talk by G. Tonelli for the CMS collaboration, 13/12/2011.
 - [37] G. Belanger, F. Boudjema, A. Pukhov, and A. Semenov, *Comput.Phys.Comm.* **176**, 367 (2007), hep-ph/0607059.
 - [38] G. Belanger, F. Boudjema, A. Pukhov, and A. Semenov, *Comput.Phys.Comm.* **180**, 747 (2009), 0803.2360.
 - [39] M. Ibe, H. Murayama, and T. Yanagida, *Phys.Rev.* **D79**, 095009 (2009), 0812.0072.
 - [40] M. Ackermann, M. Ajello, A. Albert, W. B. Atwood, L. Baldini, J. Ballet, G. Barbiellini, D. Bastieri, K. Bechtol, R. Bellazzini, et al., *PRL* **107**, 241302 (2011), 1108.3546.
 - [41] T. Cohen, J. Kearney, A. Pierce, and D. Tucker-Smith (2011), 1109.2604.
 - [42] J. Edsjo and P. Gondolo, *Phys.Rev.* **D56**, 1879 (1997), hep-ph/9704361.
 - [43] E. Aprile, K. Arisaka, F. Arneodo, A. Askin, L. Baudis, A. Behrens, K. Bokeloh, E. Brown, T. Bruch, G. Bruno, et al., *Physical Review Letters* **107**, 131302 (2011), 1104.2549.
 - [44] CDMS II Collaboration, Z. Ahmed, et al., *Science* **327**, 1619 (2010), 0912.3592.
 - [45] Z. Ahmed, D. S. Akerib, et al., *Physical Review Letters* **106**, 131302 (2011), 1011.2482.
 - [46] EDELWEISS Collaboration, E. Armengaud, et al., *Physics Letters B* **702**, 329 (2011), 1103.4070.
 - [47] M. S. Boucenna and S. Profumo, *PRD* **84**, 055011 (2011), 1106.3368.
 - [48] Choosing, for simplicity, a null mixing between S_1 and S_2 .